

Journal Pre-proof

Contributions of emotion regulation and brain structure and function to adolescent internalizing problems and stress vulnerability during the COVID-19 pandemic: A longitudinal study

David G. Weissman, Alexandra M. Rodman, Maya L. Rosen, Steven Kasperek, Makeda Mayes, Margaret A. Sheridan, Lilliana Lengua, Andrew N. Meltzoff, Katie A. McLaughlin

PII: S2667-1743(21)00047-1

DOI: <https://doi.org/10.1016/j.bpsgos.2021.06.001>

Reference: BPSGOS 27

To appear in: *Biological Psychiatry Global Open Science*

Received Date: 2 March 2021

Revised Date: 2 June 2021

Accepted Date: 3 June 2021

Please cite this article as: Weissman D.G., Rodman A.M., Rosen M.L., Kasperek S., Mayes M., Sheridan M.A., Lengua L., Meltzoff A.N. & McLaughlin K.A., Contributions of emotion regulation and brain structure and function to adolescent internalizing problems and stress vulnerability during the COVID-19 pandemic: A longitudinal study, *Biological Psychiatry Global Open Science* (2021), doi: <https://doi.org/10.1016/j.bpsgos.2021.06.001>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Inc. on behalf of Society of Biological Psychiatry.



Contributions of emotion regulation and brain structure and function to adolescent internalizing problems and stress vulnerability during the COVID-19 pandemic: A longitudinal study

David G. Weissman¹, Alexandra M. Rodman¹, Maya L. Rosen¹, Steven Kasparek¹, Makeda Mayes², Margaret A. Sheridan³, Lilliana Lengua², Andrew N. Meltzoff² & Katie A. McLaughlin¹

¹Department of Psychology, Harvard University

²Department of Psychology, University of Washington-Seattle

³Department of Psychology and Neuroscience, University of North Carolina-Chapel Hill

Corresponding Author: David G. Weissman, William James Hall, 33 Kirkland St. Cambridge, MA 02138. dweissman@fas.harvard.edu

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

Abstract

Background: Adolescence is a period of increased vulnerability for internalizing problems, particularly following stressful life events. We examine how emotion regulation and brain structure and function are associated with internalizing problems during the COVID-19 pandemic, and moderate the association between pandemic-related stressors and internalizing problems.

Methods: Data are from a longitudinal sample (N=145, aged 10-15) strategically assessed at three crucial timepoints: prior to the pandemic, early during the stay-at-home order period, and again six months later. We examined associations of amygdala and hippocampal volume and amygdala activation during an emotional processing task prior to the pandemic; use of emotion regulation strategies prior to and during the pandemic; and pandemic-related stressors with internalizing problems.

Results: Greater exposure to pandemic-related stressors was associated with higher internalizing problems both early and later in the pandemic. Youths who reported more frequent use of rumination before the pandemic and higher use of expressive suppression and lower use of cognitive reappraisal early in the pandemic had higher internalizing problems early in the pandemic. Higher left amygdala activation to neutral relative to fearful faces before the pandemic was associated with greater internalizing problems and a stronger link between pandemic-related stressors and internalizing problems early in the pandemic.

Conclusion: Pandemic-related stressors are strongly associated with adolescent internalizing problems, as are individual differences in emotional reactivity and regulation and their underlying neural mechanisms. Interventions that reduce pandemic-related stressors and foster

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

adaptive emotion regulation skills may protect against adolescent psychopathology during this period of heightened exposure to stress.

Journal Pre-proof

Introduction

Adolescence is a period of increased risk for internalizing problems (1,2). Exposure to stressful life events increases during adolescence, and adolescents experience heightened vulnerability to developing stress-related anxiety and depression symptoms (3–6). The COVID-19 pandemic has produced dramatic societal changes that have resulted in increased exposure to numerous health, economic, and social stressors for adolescents. Internalizing problems have increased during the pandemic in both adolescents and adults (7–11). However, the degree of exposure to pandemic-related stressors varies widely (10,12,13), and risk for internalizing problems related to these stressors is unlikely to be uniform. Identifying characteristics that convey risk for or resilience to internalizing problems during the COVID-19 pandemic may help to generate targets for interventions to promote well-being during this period of heightened stress exposure. This longitudinal study investigates how individual differences in multiple domains of emotional processing—use of specific emotion regulation strategies, brain structure, and neural function during emotional processing—predict vulnerability to stress-related internalizing problems during the COVID-19 pandemic.

Difficulties with emotion regulation, increased amygdala reactivity to threat, and smaller hippocampal and amygdala volume are among the psychological and neural mechanisms that may increase vulnerability for internalizing problems in children and adolescents in response to stressful life events (14–16). Use of maladaptive emotion regulation strategies—like rumination and suppression—is associated with elevated risk for internalizing problems in longitudinal studies and meta-analyses of adults and adolescents (17–20). In contrast, greater use of adaptive emotion regulation strategies, such as cognitive reappraisal (21), is less consistently associated with psychopathology than maladaptive strategies (22–24). However, more habitual and

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

effective use of cognitive reappraisal has a protective moderating influence on internalizing problems among youth who experience stressful life events (25,26). In the context of the COVID-19 pandemic, adolescents who engage in more cognitive reappraisal might be expected to be more resilient to pandemic-related stressors, while adolescents who engage in more expressive suppression and rumination might be expected to be more vulnerable to stress-related increases in internalizing problems.

Heightened amygdala reactivity to threat-related cues, such as fearful or angry faces, or scenes depicting violence, may reflect that potential threats have greater emotional salience, leading to more intense negative emotional experiences and greater mobilization of defensive responses (27–29). Heightened amygdala reactivity to potential threats is associated with depression and anxiety in adolescence, including in prospective studies (30,31). Moreover, the prospective association between stressful life events and depression is stronger among youth with higher amygdala reactivity to fearful and angry faces (15). Like the amygdala, the hippocampus is involved in the processing of emotionally salient stimuli and in regulating physiological responses to stress and negative emotion (32–34), in addition to the central role it plays in learning and memory (35). While the association between hippocampal and amygdala volume and internalizing problems are inconsistent in children and adolescents (36–40), several studies have found that lower hippocampal volume is associated with greater vulnerability to internalizing problems following stressful life events. For example, maternal aggression is associated with greater increases in depressive symptoms over time in early adolescents with smaller hippocampal volume (41), and stressful life events are associated with elevated anxiety only among participants with small hippocampal volume (42). Finally, the prospective association between stressful life events and depression symptoms is stronger among youth with

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

reduced hippocampal and amygdala volume (16). In the context of the COVID-19 pandemic, adolescents with greater amygdala reactivity to threat cues and smaller amygdala and hippocampal volume prior to the pandemic might be expected to be more vulnerable to stress-related increases in internalizing problems.

The current longitudinal study examines whether emotion regulation strategy use, amygdala reactivity to threat, and hippocampal and amygdala volume contribute to vulnerability to internalizing problems in response to COVID-19 pandemic-related stressors. We expected that greater use of rumination and expressive suppression and less frequent use of cognitive reappraisal would be associated with more internalizing problems and magnify the association between pandemic-related stressors and internalizing problems. Similarly, we expected that greater amygdala reactivity to threat and smaller amygdala and hippocampal volume—measured prior to the pandemic—would be associated with greater internalizing problems and vulnerability to stress-related internalizing problems. Due to the unprecedented nature and duration of the COVID-19 pandemic, we did not have specific hypotheses about how associations might vary between early and later in the pandemic. However, pre-pandemic internalizing problems were measured over two years earlier, and with a different scale than during the pandemic. We therefore anticipated greater stability in internalizing problems from early to later in the pandemic than from pre-pandemic to early in the pandemic. Thus, statistically, risk and protective factors would likely be more strongly associated with changes in internalizing problems early in the pandemic than later.

Methods

Participants

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

Participants were drawn from a longitudinal study of children followed from age 3 and a caregiver, with the initial aim of studying the development of self-regulation in childhood (43). At age 11-12 years, between June 2017 and October 2018, participants completed a pre-pandemic baseline as part of a study examining mechanisms linking adverse childhood experiences with psychopathology (44). Participants were excluded from the study based on the following criteria: IQ < 80, active substance dependence, psychosis, presence of pervasive developmental disorders (e.g., autism), and psychotropic medication use. We conducted two assessments of this sample during the COVID-19 pandemic. Youth and a caregiver completed questionnaires online between April and May of 2020 (Wave 1), and between November 2020 and January 2021 (Wave 2). See Figure 1 for a flowchart of participant recruitment and inclusion. Participant race and ethnicity is summarized in Table S1 in supplemental materials. Participants who had complete data at each wave did not differ from participants missing data on any variable included in study analyses.

Measures

See Supplemental Materials for greater details on all measures, including psychometrics.

Rumination (Pre-pandemic baseline). The 13-item rumination subscale of the Children's Response Styles Questionnaire (CRSQ) (45) was used to assess use of rumination.

Pandemic-related Stressors (Wave 1 and Wave 2). We developed a set of questions to assess pandemic-related stressors that was administered to children and caregivers. Stressors were coded as present if they were endorsed by either the youth or their parent and summed, using a cumulative risk approach (46). See supplemental materials and Table S2 for details. The complete scale can be found at <https://osf.io/drqku/>.

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

Cognitive Reappraisal and Expressive Suppression (Wave 1). The Emotion Regulation Questionnaire (ERQ) (47) is a 10-item self-report questionnaire assessing the tendency to use cognitive reappraisal and expressive suppression. Six items pertain to cognitive reappraisal and four pertain to expressive suppression.

Internalizing problems (Pre-pandemic baseline, Wave 1, Wave 2). Internalizing problems at the pre-pandemic baseline were assessed based on child report on the Youth Self-Report (48). At the Wave 1 and Wave 2 COVID-19 follow-up assessments adolescents completed the Strengths and Difficulties Questionnaire, a 25-item scale comprised of five subscales, with five items each (50). To make the scales comparable, internalizing scores were scaled to values between 0 and 1 (i.e. proportion of maximum scaling) (51).

Emotional Processing fMRI Task (Pre-pandemic baseline)

The emotional processing task consisted of 2 runs of 9 18-second blocks, during which participants passively viewed neutral, fearful, and scrambled face stimuli. Faces were drawn from the NimStim stimulus set (52). The “calm” faces from this dataset were used as neutral expressions, as these expressions are potentially less emotionally evocative than neutral faces (52), which can be perceived as negatively-valenced (52). Further details can be found in Supplemental Materials.

MRI Acquisition and Preprocessing

Scanning was performed on a 3T Phillips Achieva scanner at the University of Washington Integrated Brain Imaging Center (TR=2s, voxel size=3mm³ for the functional scan). Preprocessing and statistical analysis of fMRI data was performed for the purposes of extracting activation to fearful vs. neutral faces from left and right amygdala ROIs. Further details can be found in Supplemental Materials.

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

Structural MRI processing

Measures of hippocampal and amygdala volume and total intracranial volume were obtained using automatic segmentation in FreeSurfer 5.3. Following prior work (16), right and left volumes were summed to create bilateral hippocampal and amygdala volume measures. To keep variables on a similar scale for regression analyses, subcortical volumes were divided by 1,000, and intracranial volume was divided by 1,000,000. Further details can be found in Supplemental Materials.

Procedures

All study procedures were approved by the Institutional Review Board at Harvard University. Habitual use of rumination, internalizing problems, and measures of brain structure and function were assessed at the pre-pandemic baseline. Pandemic-related stressors, use of cognitive reappraisal and expressive suppression, and internalizing problems were assessed at each of the Wave 1 and Wave 2 follow-up assessments during the pandemic. Legal guardians provided informed consent and youths provided assent via electronic signature obtained using Qualtrics (Provo, UT). Once consent and assent was obtained for the pandemic follow-up assessments, parents and children were separately provided with a link to surveys on RedCap and asked to complete them. If children had trouble completing the surveys on their own, an experimenter called via phone or video chat and read the questions aloud to the child and recorded their responses.

Analyses

All analyses were conducted in R version 4.0.3 (54). We used a paired sample t-test to evaluate change in internalizing problems from the pandemic Wave 1 to Wave 2 follow-up. The associations between pandemic-related stressors and internalizing problems at the Wave 1 and

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

Wave 2 follow-ups were evaluated using multiple linear regression, controlling for internalizing problems at the previous assessment, sex, and age.

For each risk or protective factor, we fit 2 longitudinal path models. The first model examined the main effect of each factor on internalizing problems at Wave 1 and Wave 2 (Figure 2A). The second model examined the interaction between each risk and protective factor with exposure to pandemic-related stressors. For each model, both the risk or protective factor and pandemic-related stressors reported at Wave 1 and Wave 2 were mean-centered, and an interaction term was computed by multiplying them (Figure 2B). Models all fit the data well. Model fit statistics and full model output for all models is presented in Supplemental Materials.

Because 14 total path models were examined, False Discovery Rate correction (53) was used to correct for multiple comparisons within each domain of analysis (emotion regulation: 6 models; amygdala activation: 4 models; amygdala and hippocampal volume: 4 models). Therefore, False Discovery Rate-corrected *p-values* are indicated in all results and tables as *FDR-p*. Simple slopes analysis was used to follow up significant interactions. More detail, as well as reproducible code and output (Rmarkdown) of all analyses are provided in the Supplemental Materials, and all data and analysis code are available at <https://github.com/dgweissman/COVID>.

Results

Descriptive Statistics

Descriptive statistics and bivariate associations between all variables are provided in Table 1. See Figure S2 for the distribution of exposure to pandemic-related stressors.

Pandemic-related Stressors and Internalizing Problems

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

Internalizing problems increased significantly during the pandemic from Wave 1 to Wave 2 ($M_{diff}=1.18$, $95\% CI=0.59-1.77$, $t=5.80$, $df=116$). Adolescents who experienced more pandemic-related stressors early in the pandemic reported higher internalizing problems at the pandemic Wave 1 assessment (*unstandardized* $B=.834$, $SE=.134$, *standardized* $\beta=.437$, $t=6.19$, $df=138$, $p<.001$), controlling for age, sex, and internalizing problems at the pre-pandemic baseline. Notably, while the zero-order correlations between pre-pandemic internalizing problems and Wave 1 internalizing problems was small and non-significant ($r=.10$, $p=.215$), their association in the regression model that included pandemic-related stressors was modest and significant ($B=.209$, $SE=.070$, $\beta=.225$, $t=2.96$, $df=138$, $p=.004$). The association of pandemic-related stressors occurring between the Wave 1 and Wave 2 pandemic assessments was associated with higher internalizing problems at Wave 2 ($B=.486$, $SE=.166$, $\beta=.225$, $t=2.93$, $df=112$, $p=.004$), controlling for age, sex, and Wave 1 internalizing problems (Figure 3).

Emotion Regulation, Pandemic-related Stressors, and Internalizing Problems

Youths who reported more frequent use of rumination at baseline had higher internalizing problems at the Wave 1 pandemic assessment ($B=.007$, $SE=.003$, $\beta=.209$, $FDR-p=.036$), controlling for age, sex, and baseline symptoms, but not at the Wave 2 pandemic assessment controlling for age, sex, and Wave 1 symptoms ($B=.001$, $SE=.003$, $\beta=.021$, $FDR-p=.946$).

Greater use of expressive suppression reported at Wave 1 during the pandemic was associated with higher internalizing problems at Wave 1 ($B=.078$, $SE=.022$, $\beta=.265$, $FDR-p<.001$), controlling for age, sex, and pre-pandemic symptoms. Greater use of expressive suppression reported at Wave 1 during the was also associated with higher internalizing problems at Wave 2, controlling for age, sex, and Wave 1 symptoms, but this association was not significant after multiple comparisons correction ($B=.065$, $SE=.027$, $\beta=.172$, $FDR-p=.946$).

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

Greater use of cognitive reappraisal reported early in the pandemic was associated with higher internalizing problems at Wave 1, controlling for age, sex, and pre-pandemic symptoms ($B=-.120$, $SE=.023$, $\beta=-.374$, $FDR-p<.001$). However, greater use of cognitive reappraisal reported at Wave 1 during the pandemic was not associated with internalizing problems at Wave 2, controlling for age, sex, and Wave 1 symptoms ($B=-.015$, $SE=.033$, $\beta=-.037$, $FDR-p=.946$).

Use of rumination, cognitive reappraisal, and expressive suppression did not interact with pandemic-related stressors to predict internalizing problems at either point in the pandemic (see Table 2).

Amygdala Activation, Pandemic-related Stressors, and Internalizing Problems

Although highly heterogenous, amygdala activation to fearful faces was significantly greater than amygdala activation to neutral faces on average ($M=.239$, $SD=.897$, $p<.001$). Lower left amygdala activation to fearful relative to neutral faces prior to the pandemic was associated with greater internalizing problems in the early phase of the pandemic ($B=-.054$, $SE=.019$, $\beta=-.233$, $FDR-p=.012$), controlling for age, sex, and pre-pandemic symptoms. To determine whether this association was driven by greater reactivity to neutral faces or less reactivity fearful faces, we decomposed this contrast by examining the association of left amygdala activation to fearful vs scrambled faces and neutral vs scrambled faces with internalizing problems, neither of which were significant (Figure 4). No associations were observed for right amygdala activation (Table 2).

Left amygdala activation to fearful relative to neutral faces moderated the association of pandemic-related stressors with internalizing problems early in the pandemic ($B=-.028$, $SE=.010$, $\beta=-.200$, $FDR-p=.012$), controlling for age, sex, and pre-pandemic symptoms. Pandemic-related stressors were positively associated with internalizing problems among youth with low to mean

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

amygdala reactivity, but not among youth with high amygdala reactivity to fearful vs. neutral faces (Figure 5). No significant interactions were observed for right amygdala activation (Table 2).

Brain Structure, Pandemic-related Stressors, and Internalizing Problems

Neither amygdala volume nor hippocampal volume at the baseline visit were associated with internalizing problems at either point during the pandemic, nor did they interact with pandemic-related stressors in predicting internalizing problems at either wave (See Table 2).

Discussion

The COVID-19 pandemic has produced sudden and unprecedented changes in the lives of adolescents that have introduced many novel stressors and exacerbated risk for anxiety and depression (7–11), which were already on the rise among adolescents in the United States (54). The current study examined factors contributing to individual differences in the impact of the pandemic on adolescent internalizing problems in the early phase of the pandemic (late spring 2020) and again six months later (late fall/winter 2020). Adolescents demonstrated worsening internalizing problems over the course of the pandemic, and adolescents with high exposure to pandemic-related stressors had particularly elevated internalizing problems relative to the previous timepoint. We found significant variability in internalizing problems based on the use of different emotion regulation strategies before and during the pandemic—including rumination, cognitive reappraisal, and expressive suppression – as well as amygdala activation measured prior to the pandemic. Specifically, greater left amygdala activation to neutral relative to fearful faces was associated with higher internalizing problems during the pandemic. These findings support widespread concerns about the mental health consequences of the pandemic on adolescent mental health, particularly for those exposed to more pandemic-related stressors. Use

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

of maladaptive emotion regulation strategies and less differentiated neural responses to threatening and neutral cues may make youth particularly vulnerable to the increased exposure to stressors that have characterized the pandemic.

The degree of exposure to pandemic-related stressors was associated with worsening internalizing problems over time both early in the pandemic and as the pandemic progressed. This is consistent with cumulative risk models, which argue that stressors have a cumulative impact on mental health (46), and prior work on exposure to stressors and psychopathology during community-wide disruptions like natural disasters and terrorist attacks (55–57). These findings therefore characterize the impact of a unique and unprecedented source of stress, but conform to the general patterns of association observed in other stressful contexts. Further, although the stressors examined here (e.g., social isolation, parental job loss, food insecurity) were exacerbated by the pandemic and lockdowns, these types of stressors can and do occur in other contexts. Conversely, social isolation due to the lockdowns may have further exacerbated the mental health impacts of these stressors by limiting the social support available to adolescents. While the pandemic continues, policies aimed at mitigating the stressors experienced by families (e.g. unemployment benefits, return to in-person schooling) may have the greatest potential to decrease the impact of the pandemic on youth mental health.

Internalizing problems pre-pandemic were not correlated with internalizing problems during the pandemic. However, they were significantly associated in the regression model that included pandemic-related stressors, with an effect size consistent with the stability that might be expected between two moderately correlated measures (57) collected 2-3 years apart during adolescence, suggesting that the strong association between pandemic-related stressors and

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

internalizing problems may have reduced the stability of internalizing problems in early adolescence.

Greater use of maladaptive emotion regulation strategies—including rumination and expressive suppression— and lower use of cognitive reappraisal predicted worse internalizing problems early in the pandemic. Overall, greater use of adaptive emotion regulation strategies (i.e., cognitive reappraisal) and lower use of maladaptive strategies (i.e., rumination and suppression) was associated with lower internalizing problems during the pandemic, but was not as effective at reducing the impact of pandemic-related stressors. Further, perhaps due to the high rank-order stability of internalizing problems during the pandemic, the protective influence of emotion regulation strategy use was specific to the early months of the pandemic. Nonetheless, these results provide some evidence that interventions focused on teaching adolescents adaptive emotion regulation skills may be an effective means of reducing internalizing psychopathology, during the pandemic and beyond.

Higher left amygdala activation to neutral compared to fearful faces prior to the pandemic had a modest and significant association with internalizing problems in the early phase of the pandemic and moderated the association of pandemic-related stressor with internalizing problems, such that the association was stronger among youth with higher left amygdala activation to neutral relative to fearful faces. The direction of these associations is unexpected. In prior work, the association between stressful life events and depression was stronger in youth with greater amygdala responses to fearful and angry faces compared to shapes (15). Decomposing our finding by examining fearful and neutral faces each compared to scrambled faces was inconclusive. However, our results may nonetheless reflect that youth who differentiate less between fearful faces—which indicate the presence of a potential threat in the

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

environment, and neutral faces—which are ambiguous and do not clearly reflect the presence of a threat, may interpret neutral or ambiguous cues as more threatening. Amygdala reactivity to potential threats is amplified in contexts of high uncertainty (59,60). Youth who experience violence and other forms of childhood trauma—experiences characterized by high threat and uncertainty that confer greater risk for internalizing problems—are more likely to perceive neutral faces as angry (61,62) and to attribute hostile intent to ambiguous social cues more generally (63). Less differentiated amygdala reactivity to threatening compared to neutral cues, perhaps reflecting greater uncertainty in the interpretation of the more ambiguous neutral faces, may make adolescents more vulnerable to developing internalizing problems when faced with chronic stress and uncertainty, common experiences during the COVID-19 pandemic. Indeed, amygdala activation was not correlated with internalizing problems assessed concurrently at the pre-pandemic baseline, which suggests that less differentiated amygdala reactivity to threatening compared to neutral cues may be more an indicator of vulnerability to future stress and uncertainty than a direct correlate of internalizing problems.

Neither amygdala nor hippocampal volume was associated with internalizing problems directly nor did they interact with pandemic-related stressors in predicting internalizing problems at either time point in the pandemic. Thus, we failed to replicate the pattern of smaller hippocampal and amygdala volume being associated with greater vulnerability to stress that has been observed previously among youth exposed to childhood violence (16,42). It is possible that differences in hippocampal and amygdala volume resulting from exposure to traumatic stressors may reflect specific structural alterations that impact responses to stress more so than the variability in hippocampal and amygdala volume observed in samples without high trauma exposure.

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

This study confirms and replicates the concerning impact that the pandemic has had on adolescent mental health (7–11) and reveals several sources of meaningful variability in symptoms of depression and anxiety among adolescents. Exposure to stressors is by far the most robust and consistent predictor of internalizing problems examined in this study, consistent with substantial evidence on the link between stressors and internalizing psychopathology (64,65). Adolescent-reported use of maladaptive emotion regulation strategies was also a modest predictor that can be measured inexpensively and easily and could help to identify youths who might be particularly at risk for developing internalizing problems and thus benefit from mental health interventions. Alternatively, while the modest associations of lower amygdala discrimination between fearful and neutral faces in increasing vulnerability to pandemic-related stressors informs our understanding of the mechanisms underlying stress sensitivity in adolescents, this measure is substantially more costly and difficult to acquire, and may offer marginal additional clinical or practical utility in identifying the most vulnerable youths relative to self- and parent-reports of stressors and emotion regulation strategy use.

While this study has several strengths, including rich, multi-informant (parent and adolescent) and multimodal (self-reports, neuroimaging) data on a relatively large sample of adolescents assessed at three time points before and during the COVID-19 pandemic, it also has several limitations that should be considered when interpreting our findings. First, because the participants in this sample were all within a relatively narrow two-year age band, we are unable to characterize the specificity or generalizability of these findings across other developmental windows. Second, the stimuli in the fMRI task of the current study included only fearful and neutral faces. Therefore, we are unable to determine whether the association between reduced differentiation in the amygdala between fearful and neutral faces and internalizing problems in

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

the context of pandemic-related stressors is specific to fear, or may reflect responses to negative emotional cues more generally. Third, because each of our neuroimaging and emotion regulation measures were collected at only one time point, we were unable to examine their stability or the consistency of their associations with internalizing problems across multiple timepoints as we were with pandemic-related stressors. Finally, while the pandemic-related stress measure demonstrated convergent validity via consistently moderate associations with internalizing problems at both waves, it nonetheless had the limitation of equally weighting stressors that may have had highly variable objective and subjective impacts. Despite these limitations, when synthesized with the many other developmental studies underway during the pandemic, we anticipate these longitudinal, multi-informant, multimodal findings contributing to a richer and more complete understanding of factors contributing to adolescent mental health during the pandemic, and factors contributing to stress vulnerability more broadly.

Internalizing problems have increased steadily among adolescents during the COVID-19 pandemic, particularly for those who have experienced a high number of pandemic-related stressors, as well among adolescents who habitually use maladaptive emotion regulation strategies like rumination and suppression and use cognitive reappraisal less frequently. Lower differentiation of amygdala activation between threatening and neutral stimuli, assessed prior to the pandemic, is also associated with increased vulnerability to stress-related internalizing problems. Attempts to identify adolescents most in need of mental health interventions would benefit from screening for exposure to pandemic-related stressful experiences and use of emotion regulation strategies. Such efforts may help to identify adolescents in need of extra support and intervention to mitigate the mental health crisis posed by the COVID-19 pandemic.

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

Acknowledgements

The initial conceptualization of the current study was done by KAM and ANM. This work was supported by the Bezos Family Foundation (to ANM) for collection of data with additional support by the National Institute of Mental Health (R01 MH106482 to KAM) and the National Institute of Child Development (R01 HD054465 to LL). The authors would also like to acknowledge Reshma Sreekala and Malila Freeman for help with data collection and Frances Li for help with compiling surveys.

Financial Disclosures

The authors report no biomedical financial interests or potential conflicts of interest.

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

References

1. Hankin BL, Abramson LY, Moffitt TE, Silva PA, McGee R, Angell KE (1998): Development of depression from preadolescence to young adulthood: emerging gender differences in a 10-year longitudinal study. *J Abnorm Psychol* 107: 128–140.
2. Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE (2005): Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 62: 593–602.
3. La Greca AM, Harrison HM (2005): Adolescent Peer Relations, Friendships, and Romantic Relationships: Do They Predict Social Anxiety and Depression? *J Clin Child Adolesc Psychol* 34: 49–61.
4. Siegel RS, La Greca AM, Harrison HM (2009): Peer Victimization and Social Anxiety in Adolescents: Prospective and Reciprocal Relationships. *J Youth Adolesc* 38: 1096–1109.
5. Larson R, Ham M (1993): Stress and “Storm and Stress” in Early Adolescence: The Relationship of Negative Events with Dysphoric Affect. *Dev Psychol* 29: 130–40.
6. Larson R, Lampman-Petratis C (1989): Daily Emotional States as Reported by Children and Adolescents. *Child Dev* 60: 1250–1260.
7. Chahal R, Kirshenbaum JS, Miller JG, Ho TC, Gotlib IH (2020): Higher executive control network coherence buffers against puberty-related increases in internalizing symptoms during the COVID-19 pandemic. *Biol Psychiatry Cogn Neurosci Neuroimaging*.
<https://doi.org/10.1016/j.bpsc.2020.08.010>
8. Crescentini C, Feruglio S, Matiz A, Paschetto A, Vidal E, Cogo P, Fabbro F (2020): Stuck Outside and Inside: An Exploratory Study on the Effects of the COVID-19 Outbreak on

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

- Italian Parents and Children's Internalizing Symptoms. *Front Psychol* 11.
<https://doi.org/10.3389/fpsyg.2020.586074>
9. Loades ME, Chatburn E, Higson-Sweeney N, Reynolds S, Shafran R, Brigden A, *et al.* (2020): Rapid Systematic Review: The Impact of Social Isolation and Loneliness on the Mental Health of Children and Adolescents in the Context of COVID-19. *J Am Acad Child Adolesc Psychiatry*. <https://doi.org/10.1016/j.jaac.2020.05.009>
10. Whittle S, Bray K, Lin S, Schwartz O (2020): *Parenting and Child and Adolescent Mental Health during the COVID-19 Pandemic*. PsyArXiv. <https://doi.org/10.31234/osf.io/ag2r7>
11. Racine N, Cooke JE, Eirich R, Korczak DJ, McArthur B, Madigan S (2020): Child and adolescent mental illness during COVID-19: A rapid review. *Psychiatry Res* 292: 113307.
12. McLaren J (2020): *Racial Disparity in COVID-19 Deaths: Seeking Economic Roots with Census Data*. National Bureau of Economic Research. <https://doi.org/10.3386/w27407>
13. Montenovo L, Jiang X, Rojas FL, Schmutte IM, Simon KI, Weinberg BA, Wing C (2020): *Determinants of Disparities in Covid-19 Job Losses*. National Bureau of Economic Research. <https://doi.org/10.3386/w27132>
14. McLaughlin KA, Koenen KC, Bromet EJ, Karam EG, Liu H, Petukhova M, *et al.* (2017): Childhood adversities and post-traumatic stress disorder: Evidence for stress sensitisation in the World Mental Health Surveys. *Br J Psychiatry*.
<https://doi.org/10.1192/bjp.bp.116.197640>
15. Swartz JR, Knodt AR, Radtke SR, Hariri AR (2015): A Neural Biomarker of Psychological Vulnerability to Future Life Stress. *Neuron* 85: 505–511.

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

16. Weissman DG, Lambert HK, Rodman AM, Peverill M, Sheridan MA, McLaughlin KA (2020): Reduced hippocampal and amygdala volume as a mechanism underlying stress sensitization to depression following childhood trauma. *Depress Anxiety*.
<https://doi.org/10.1002/da.23062>
17. Aldao A, Nolen-Hoeksema S, Schweizer S (2010): Emotion-regulation strategies across psychopathology: A meta-analytic review. *Clin Psychol Rev*.
<https://doi.org/10.1016/j.cpr.2009.11.004>
18. Nolen-Hoeksema S, Aldao A (2011): Gender and age differences in emotion regulation strategies and their relationship to depressive symptoms. *Personal Individ Differ*.
<https://doi.org/10.1016/j.paid.2011.06.012>
19. McLaughlin KA, Hatzenbuehler ML, Mennin DS, Nolen-Hoeksema S (2011): Emotion dysregulation and adolescent psychopathology: A prospective study. *Behav Res Ther*.
<https://doi.org/10.1016/j.brat.2011.06.003>
20. McLaughlin KA, Nolen-Hoeksema S (2011): Rumination as a transdiagnostic factor in depression and anxiety. *Behav Res Ther*. <https://doi.org/10.1016/j.brat.2010.12.006>
21. Gross JJ, John OP (2003): Individual Differences in Two Emotion Regulation Processes: Implications for Affect, Relationships, and Well-Being. *J Pers Soc Psychol*.
<https://doi.org/10.1037/0022-3514.85.2.348>
22. Aldao A, Nolen-Hoeksema S (2012): The influence of context on the implementation of adaptive emotion regulation strategies. *Behav Res Ther*.
<https://doi.org/10.1016/j.brat.2012.04.004>

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

23. Aldao A, Nolen-Hoeksema S, Schweizer S (2010): Emotion-regulation strategies across psychopathology: A meta-analytic review. *Clinical Psychology Review*.
<https://doi.org/10.1016/j.cpr.2009.11.004>
24. Aldao A, Nolen-Hoeksema S (2012): When are adaptive strategies most predictive of psychopathology? *J Abnorm Psychol*. <https://doi.org/10.1037/a0023598>
25. Flouri E, Mavroveli S (2013): Adverse Life Events and Emotional and Behavioural Problems in Adolescence: The Role of Coping and Emotion Regulation. *Stress Health* 29: 360–368.
26. Rodman AM, Jenness JL, Weissman DG, Pine DS, McLaughlin KA (2019): Neurobiological Markers of Resilience to Depression Following Childhood Maltreatment: The Role of Neural Circuits Supporting the Cognitive Control of Emotion. *Biol Psychiatry* 86: 464–473.
27. Costafreda SG, Brammer MJ, David AS, Fu CHY (2008): Predictors of amygdala activation during the processing of emotional stimuli: A meta-analysis of 385 PET and fMRI studies. *Brain Res Rev*. <https://doi.org/10.1016/j.brainresrev.2007.10.012>
28. Cunningham WA, Brosch T (2012): Motivational Salience: Amygdala Tuning From Traits, Needs, Values, and Goals. *Curr Dir Psychol Sci* 21: 54–59.
29. Adolphs R, Tranel D, Damasio H, Damasio A (1995): Fear and the human amygdala. *J Neurosci* 15.
30. Swartz JR, Williamson DE, Hariri AR (2015): Developmental change in amygdala reactivity during adolescence: Effects of family history of depression and stressful life events. *Am J Psychiatry* 172: 276–283.

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

31. Pagliaccio D, Luby JL, Luking KR, Belden AC, Barch DM (2014): Brain-behavior relationships in the experience and regulation of negative emotion in healthy children: Implications for risk for childhood depression. *Dev Psychopathol.*
<https://doi.org/10.1017/S0954579414001035>
32. Frodl T, O'Keane V (2013): How does the brain deal with cumulative stress? A review with focus on developmental stress, HPA axis function and hippocampal structure in humans. *Neurobiol Dis.* <https://doi.org/10.1016/j.nbd.2012.03.012>
33. Gläscher J, Adolphs R, Flannery J, Goff B, Telzer EH, Shapiro M, *et al.* (2003): Processing of the arousal of subliminal and supraliminal emotional stimuli by the human amygdala. *J Neurosci Off J Soc Neurosci* 23: 10274–82.
34. Williams LM, Das P, Liddell B, Olivieri G, Peduto A, Brammer MJ, Gordon E (2005): BOLD, sweat and fears: fMRI and skin conductance distinguish facial fear signals. *NeuroReport* 16: 49–52.
35. Lupien SJ, Lepage M (2001): Stress, memory, and the hippocampus: can't live with it, can't live without it. *Behav Brain Res* 127: 137–158.
36. Caetano SC, Fonseca M, Hatch JP, Olvera RL, Nicoletti M, Hunter K, *et al.* (2007): Medial temporal lobe abnormalities in pediatric unipolar depression. *Neurosci Lett.*
<https://doi.org/10.1016/j.neulet.2007.06.014>
37. Ellis R, Fernandes A, Simmons JG, Mundy L, Patton G, Allen NB, Whittle S (2019): Relationships between adrenarcheal hormones, hippocampal volumes and depressive symptoms in children. *Psychoneuroendocrinology.*
<https://doi.org/10.1016/j.psyneuen.2019.02.016>

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

38. MacMaster FP, Kusumakar V (2004): Hippocampal volume in early onset depression. *BMC Med.* <https://doi.org/10.1186/1741-7015-2-2>
39. MacMillan S, Szeszko PR, Moore GJ, Madden R, Lorch E, Ivey J, *et al.* (2003): Increased Amygdala: Hippocampal Volume Ratios Associated with Severity of Anxiety in Pediatric Major Depression. *J Child Adolesc Psychopharmacol.* <https://doi.org/10.1089/104454603321666207>
40. Rosso IM, Cintron CM, Steingard RJ, Renshaw PF, Young AD, Yurgelun-Todd DA (2005): Amygdala and hippocampus volumes in pediatric major depression. *Biol Psychiatry.* <https://doi.org/10.1016/j.biopsych.2004.10.027>
41. Whittle S, Yap MBH, Sheeber L, Dudgeon P, Yücel M, Pantelis C, *et al.* (2011): Hippocampal volume and sensitivity to maternal aggressive behavior: A prospective study of adolescent depressive symptoms. *Development and Psychopathology.* <https://doi.org/10.1017/S0954579410000684>
42. Gorka AX, Hanson JL, Radtke SR, Hariri AR (2014): Reduced hippocampal and medial prefrontal gray matter mediate the association between reported childhood maltreatment and trait anxiety in adulthood and predict sensitivity to future life stress. *Biol Mood Anxiety Disord.* <https://doi.org/10.1186/2045-5380-4-12>
43. Lengua LJ, Moran L, Zalewski M, Ruberry E, Kiff C, Thompson S (2015): Relations of Growth in Effortful Control to Family Income, Cumulative Risk, and Adjustment in Preschool-age Children. *J Abnorm Child Psychol* 43: 705–720.
44. Cuartas J, Weissman DG, Sheridan MA, Lengua L, McLaughlin KA (2021): Corporal Punishment and Elevated Neural Response to Threat in Children. *Child Dev.* <https://doi.org/10.1111/cdev.13565>

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

45. Abela JRZ, Brozina K, Haigh EP (2002): An examination of the response styles theory of depression in third- and seventh-grade children: A short-term longitudinal study. *J Abnorm Child Psychol*. <https://doi.org/10.1023/A:1019873015594>
46. Evans GW, Li D, Whipple SS (2013): Cumulative risk and child development. *Psychol Bull* 139: 1342–1396.
47. Gross JJ, John OP (2003): Individual Differences in Two Emotion Regulation Processes: Implications for Affect, Relationships, and Well-Being. *J Pers Soc Psychol*. <https://doi.org/10.1037/0022-3514.85.2.348>
48. Achenbach TM (1991): Manual for the child behavior checklist/4-18 and 1991 profile. Burlington VT.
49. Goodman R (1997): The Strengths and Difficulties Questionnaire. *J Child Psychol Psychiatry* 38: 581–586.
50. Goodman A, Lamping DL, Ploubidis GB (2010): When to Use Broader Internalising and Externalising Subscales Instead of the Hypothesised Five Subscales on the Strengths and Difficulties Questionnaire (SDQ): Data from British Parents, Teachers and Children. *J Abnorm Child Psychol* 38: 1179–1191.
51. Little TD (2013): *Longitudinal Structural Equation Modeling*. Guilford Press.
52. Tottenham N, Tanaka JW, Leon AC, McCarry T, Nurse M, Hare TA, *et al.* (2009): The NimStim set of facial expressions: Judgments from untrained research participants. *Psychiatry Res* 168: 242–249.
53. Benjamini Y, Hochberg Y (1995): Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J R Stat Soc Ser B Methodol* 57: 289–300.

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

54. Keyes KM, Gary D, O'Malley PM, Hamilton A, Schulenberg J (2019): Recent increases in depressive symptoms among US adolescents: trends from 1991 to 2018. *Soc Psychiatry Psychiatr Epidemiol* 54: 987–996.
55. Mclaughlin KA, Fairbank JA, Gruber MJ, Jones RT, Lakoma MD, Pfefferbaum B, *et al.* (2009): Serious emotional disturbance among youths exposed to Hurricane Katrina 2 years postdisaster. *J Am Acad Child Adolesc Psychiatry* 48: 1069–1078.
56. Fairbrother G, Stuber J, Galea S, Fleischman AR, Pfefferbaum B (2003): Posttraumatic stress reactions in new York City children after the September 11, 2001, terrorist attacks. *Ambul Pediatr Off J Ambul Pediatr Assoc* 3: 304–311.
57. Lengua LJ, Long AC, Meltzoff AN (2006): Pre-attack stress-load, appraisals, and coping in children's responses to the 9/11 terrorist attacks. *J Child Psychol Psychiatry* 47: 1219–1227.
58. Theunissen MHC, de Wolff MS, Reijneveld SA (2019): The Strengths and Difficulties Questionnaire Self-Report: A Valid Instrument for the Identification of Emotional and Behavioral Problems. *Acad Pediatr* 19: 471–476.
59. Hsu M, Bhatt M, Adolphs R, Tranel D, Camerer CF (2005): Neural Systems Responding to Degrees of Uncertainty in Human Decision-Making. *Science* 310: 1680–1683.
60. Sarinopoulos I, Grupe DW, Mackiewicz KL, Herrington JD, Lor M, Steege EE, Nitschke JB (2010): Uncertainty during Anticipation Modulates Neural Responses to Aversion in Human Insula and Amygdala. *Cereb Cortex N Y NY* 20: 929–940.
61. Pollak SD, Messner M, Kistler DJ, Cohn JF (2009): Development of Perceptual Expertise in Emotion Recognition. *Cognition* 110: 242–247.

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

62. Pollak SD, Cicchetti D, Hornung K, Reed A (2000): Recognizing emotion in faces: developmental effects of child abuse and neglect. *Dev Psychol*.
<https://doi.org/10.1037/0012-1649.36.5.679>
63. Crick NR, Dodge KA (1994): Review and reformulation of social information-processing mechanisms in children's social adjustment. *Psychol Bull* 115: 74–101.
64. Grant KE, Compas BE, Thurm AE, McMahon SD, Gipson PY (2004): Stressors and child and adolescent psychopathology: measurement issues and prospective effects. *J Clin Child Adolesc Psychol Off J Soc Clin Child Adolesc Psychol Am Psychol Assoc Div 53* 33: 412–425.
65. Grant KE, Compas BE, Stuhlmacher AF, Thurm AE, McMahon SD, Halpert JA (2003): Stressors and child and adolescent psychopathology: moving from markers to mechanisms of risk. *Psychol Bull* 129: 447–466.

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

Table 1: Descriptive statistics and bivariate correlations

	Baseline			Pandemic Follow-up 1														
	<i>N</i>	<i>M</i>	<i>SD</i>	<i>N</i>	<i>M</i>	<i>SD</i>												
1. Sex (Female)	145	.43	.50	10. Age	143	14.4	.458											
2. Income-to-needs ratio	143	3.75	1.79	11. Pandemic-related stressors	143	2.28	1.74											
3. YSR Internalizing Problems	145	51.7	10.8	12. SDQ Internalizing problems	143	3.79	3.32											
4. Rumination	145	7.88	6.21	13. Cognitive reappraisal	143	3.26	.647											
5. Intracranial volume†	122	1.55	.145	14. Expressive suppression	143	2.84	.706											
6. Amygdala Volume‡	122	3.08	.357	Pandemic Follow-up 2														
7. Hippocampal volume‡	122	8.35	.780	15. Age	124	14.9	.477											
8. Left Amygdala activation	122	.239	.897	16. Pandemic-related stressors	124	2.16	1.71											
9. Right Amygdala activation	122	.158	.867	17. SDQ Internalizing problems	124	5.05	3.75											
Correlations																		
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	
1	-																	
2	.21*	-																
3	-.38*	-.26*	-															
4	-.16	.01	.50*	-														
5	-.52*	.11	.09	-.00	-													
6	-.50*	.17	.04	-.00	.66*	-												
7	-.46*	.22*	.10	-.01	.64*	.69*	-											
8	-.11	.00	.09	-.02	-.01	-.02	-.09	-										
9	.01	-.02	.07	-.04	-.07	-.05	-.08	.57*	-									
10	.05	.42*	-.12	.02	.02	.01	-.04	.06	-.05	-								
11	.04	-.13	.00	.03	-.04	-.12	-.02	-.26*	-.26*	-.05	-							
12	.28*	-.14	.10	.21*	-.31*	-.29*	-.30*	-.25*	-.03	.00	.46*	-						
13	.00	.10	-.11	.00	.00	.00	.01	.12	-.08	-.04	-.26*	-.39*	-					
14	.08	-.08	.05	.03	-.12	-.13	-.14	-.04	-.02	.10	.10	.30*	-.16	-				
15	.03	.40*	-.18	.02	.03	.01	.02	-.05	-.12	1.00*	.07	.11	-.08	.10	-			
16	.07	-.25*	.09	.02	-.20*	-.40*	-.39*	-.14	-.15	-.12	.42*	.30*	-.09	.15	-.08	-		
17	.26*	.02	.07	.13	-.30*	-.26*	-.24*	-.14	.03	-.04	.26*	.59*	-.23*	.32*	-.03	.37*	-	

Note: * $p < .05$, YSR = Youth Self-report, SDQ = Strength and Difficulties Questionnaire.. † units = $\text{mm}^3 \times 10^6$, ‡ units = $\text{mm}^3 \times 10^3$

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

Table 2: Results of Regression analyses from Longitudinal Path Models

Independent Variables	Wave 1 Internalizing Problems					Wave 2 Internalizing Problems				
	<i>B</i>	<i>SE</i>	β	<i>p</i>	<i>FDR-p</i>	<i>B</i>	<i>SE</i>	β	<i>p</i>	<i>FDR-p</i>
Emotion Regulation										
Rumination	.007	.003	.209	.018	.036	.001	.003	.021	.791	.946
Expressive Suppression	.078	.022	.265	.000	.000	.065	.027	.172	.018	.108
Cognitive Reappraisal	-.120	.023	-.374	.000	.000	-.015	.033	-.037	.647	.946
Rumination x Stress	.000	.001	-.013	.842	.842	-.001	.002	-.046	.537	.946
Expressive Suppression x Stress	.023	.011	.136	.044	.066	.001	.014	.005	.946	.946
Cognitive Reappraisal x Stress	-.011	.011	-.067	.335	.402	-.003	.012	-.017	.822	.946
Amygdala activation										
Left Amygdala	-.054	.019	-.233	.005	.012	-.005	.023	-.017	.830	.939
Right Amygdala	-.008	.021	-.034	.694	.802	-.006	.024	-.019	.811	.939
Left Amygdala Reactivity x Stress	-.028	.010	-.200	.006	.012	-.001	.013	-.006	.939	.939
Right Amygdala Reactivity x Stress	-.003	.010	-.022	.789	.802	.014	.013	.083	.280	.939
Brain structure										
Amygdala Volume	-.047	.067	-.080	.478	.637	.060	.077	.079	.439	.785
Hippocampal Volume	-.027	.029	-.099	.364	.637	.019	.035	.055	.585	.785
Amygdala Volume x Stress	-.044	.025	-1.156	.083	.332	-.005	.037	-.103	.888	.888
Hippocampal Volume x Stress	-.004	.014	-.021	.784	.784	-.009	.016	-.043	.589	.785

Note: Each row represents a separate path model. Covariates for all models were age, sex, and internalizing problems at the previous wave. Full results of each model and reproducible analysis code can be found in supplemental materials. *B* = unstandardized coefficient, β = standardized coefficient, *p* = uncorrected p-values, *FDR-p* = false discovery rate corrected p-values using the Benjamini & Hochberg procedure.

PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

Figure 1: Participant Recruitment and Inclusion Flow Chart

Figure 1: Flow chart depicts recruitment and participation at each wave of data collection

Figure 2: Longitudinal Path Models

Figure 2: Diagrams represent the path models used to test study hypotheses. Covariances were freely estimated. Models testing associations with hippocampal and amygdala volume also included paths from total intracranial volume to internalizing at Wave 1 and Wave 2 and covariances with the other predictors. Pre = pre-pandemic baseline, W1 = Wave 1, W2 = Wave 2, Interaction Term = product of risk/protective factor and pandemic-related stressors (both mean-centered).

Figure 3: Association between Pandemic-Related Stressors and Internalizing Problems

Figure 3: Exposure to pandemic-related stressors is positively associated with internalizing problems at both Wave 1 (April-May 2020) and Wave 2 (November 2020-January 2021).

Figure 4: Left Amygdala Discrimination between Fearful and Neutral Faces and Internalizing Problems

Figure 4: Left amygdala reactivity to fearful vs. neutral faces was negatively associated with internalizing problems early in the pandemic. Examining the association of both fearful faces and neutral faces relative to scrambled faces revealed that this association was attributable to both greater activation to neutral faces and lower activation to fearful faces, but neither of those contrasts were significantly associated with internalizing problems.

Figure 5: Amygdala Discrimination, Pandemic-Related Stressors, and Internalizing Symptoms

Figure 5: Pandemic-related stressors are more strongly associated with internalizing problems in participants with lower amygdala reactivity to fearful vs. neutral faces (i.e., greater


PANDEMIC STRESS AND TEEN INTERNALIZING PROBLEMS

differentiation of threatening vs. neutral stimuli). Among participants with high amygdala response to fearful vs. neutral faces, the association between pandemic-related stressors and internalizing problems is not significant.

Journal Pre-proof

227 total participants with pre-pandemic questionnaire data (YSR, Rumination)


183 completed neuroimaging (Amygdala activation, hippocampal and amygdala volume)



6 excluded for image quality. 177 participants with usable data



145 completed Wave 1 assessment (Pandemic-related stress, SDQ, ERQ)

122 completed neuroimaging and Wave 1 assessment




124 completed Wave 2 assessment (Pandemic-related stress, SDQ)


102 completed neuroimaging and Wave 2 assessment





A. Main Effects Models




B. Interaction with Pandemic-Related Stress Models






Internalizing Problems



Internalizing Problems



Left Amygdala Activation (Fear vs. Neutral)

